

ABSTRACT

The present invention relates to nucleic acid molecules encoding cell surface receptors on immune cells and the characteristic peptides that comprise these receptors. More specifically, the present invention concerns the use of synthetic and recombinant peptides comprising natural killer ("NK") cell surface receptors. The synthetic and recombinant peptides are used to generate monoclonal antibodies that bind a specific NK cell surface receptor called CS1. The binding of the monoclonal antibody to the NK cell surface receptor leads to NK cell activation. In a particular embodiment of the present invention, the monoclonal antibodies are utilized in a method that inhibits the growth of tumor cells.

The following documents and publications are incorporated by reference herein.

REFERENCES CITED

U.S. PATENT DOCUMENTS

U.S. Patent No. 5,929,237 issued on July 27, 1999 with Kahn as inventor.
U.S. Patent No. 5,859,184 issued on January 12, 1999 with Kahn as inventor.
U.S. Patent No. 5,840,833 issued on November 24, 1998 with Kahn as inventor.
U.S. Patent No. 5,710,245 issued on January 20, 1998 with Kahn as inventor.
U.S. Patent No. 5,674,976 issued on October 7, 1997 with Kahn as inventor.
U.S. Patent No. 5,672,681 issued on September 30, 1997 with Kahn as inventor.
U.S. Patent No. 5,670,155 issued on September 23, 1997 with Kahn as inventor.
U.S. Patent No. 5,618,914 issued on April 8, 1997 with Kahn as inventor.
U.S. Patent No. 5,475,085 issued on December 12, 1995 with Kahn as inventor.
U.S. Patent No. 5,446,128 issued on August 29, 1995 with Kahn as inventor.
U.S. Patent No. 5,440,013 issued on August 8, 1995 with Kahn as inventor.
U.S. Patent No. 4,554,101 issued on November 19, 1985 with Hopp as inventor.

PUBLICATIONS

Coffey, A. J., Brooksbank, R. A., Brandau, O., Oohashi, T., Howell, G. R., Bye, J. M.,
Cahn, A. P., Durham, J., Heath, P., Wray, P., Pavitt, R., Wilkinson, J., Leversha, M.,
Huckle, E., Shaw-Smith, C. J., Dunham, A., Rhodes, S., Schuster, V., Porta, G., Yin,
L., Serafini, P., Sylla, B., Zollo, M., Franco, B., Bentley, D. R., and et al. (1998) Host
5 response to EBV infection in X-linked lymphoproliferative disease results from
mutations in an SH2-domain encoding gene . Nat Genet 20: 129-35

- Davis, S. J., Ikemizu, S., Wild, M. K., and van der Merwe, P. A. (1998) CD2 and the nature of protein interactions mediating cell-cell recognition. *Immunol Rev* 163: 217-36
- Kato, K., Koyanagi, M., Okada, H., Takanashi, T., Wong, Y. W., Williams, A. F., Okumura, K., and Yagita, H. (1992) CD48 is a counter-receptor for mouse CD2 and is involved in T cell activation. *J Exp Med* 176: 1241-9
- Kumaresan, P. R. and Mathew, P. A. (2000) Structure of the human natural killer cell receptor *2B4* gene and identification of a novel alternative transcript. *Immunogenetics* in press, DOI 10.1007/s002510000237:
- Mathew, P. A., Garni-Wagner, B. A., Land, K., Takashima, A., Stoneman, E., Bennett, M., and Kumar, V. (1993) Cloning and characterization of the *2B4* gene encoding a molecule associated with non-MHC-restricted killing mediated by activated natural killer cells and T cells. *J Immunol* 151: 5328-37
- Sayos, J., Wu, C., Morra, M., Wang, N., Zhang, X., Allen, D., van Schaik, S., Notarangelo, L., Geha, R., Roncarolo, M. G., Oettgen, H., De Vries, J. E., Aversa, G., and Terhorst, C. (1998) The X-linked lymphoproliferative-disease gene product SAP regulates signals induced through the co-receptor SLAM . *Nature* 395: 462-9
- Sewell, W. A., Palmer, R. W., Spurr, N. K., Sheer, D., Brown, M. H., Bell, Y., and Crumpton, M. J. (1988) The human LFA-3 gene is located at the same chromosome band as the gene for its receptor CD2. *Immunogenetics* 28: 278-82

- Smith, G. M., Biggs, J., Norris, B., Anderson-Stewart, P., and Ward, R. (1997)
 Detection of a soluble form of the leukocyte surface antigen CD48 in
 plasma and its elevation in patients with lymphoid leukemias and arthritis. *J*
Clin Immunol 17: 502-9
- 5 Stepp, S. E., Schatzle, J. D., Bennett, M., Kumar, V., and Mathew, P. A. (1999)
 Gene structure of the murine NK cell receptor 2B4: presence of two
 alternatively spliced isoforms with distinct cytoplasmic domains. *Eur J*
Immunol 29: 2392-9
- 10 Tangye, S. G., Lazetic, S., Woollatt, E., Sutherland, G. R., Lanier, L. L., and
 Phillips, J. H. (1999) Cutting edge: human 2B4, an activating NK cell receptor,
 recruits the protein tyrosine phosphatase SHP-2 and the adaptor signaling protein
 SAP. *J Immunol* 162: 6981-5
- 15 van der Merwe, P. A., Barclay, A. N., Mason, D. W., Davies, E. A., Morgan, B. P.,
 Tone, M., Krishnam, A. K., Ianelli, C., and Davis, S. J. (1994) Human cell-
 adhesion molecule CD2 binds CD58 (LFA-3) with a very low affinity and
 an extremely fast dissociation rate but does not bind CD48 or CD59.
Biochemistry 33: 10149-60
- 20 van der Merwe, P. A., McPherson, D. C., Brown, M. H., Barclay, A. N., Cyster, J.
 G., Williams, A. F., and Davis, S. J. (1993) The NH2-terminal domain of
 rat CD2 binds rat CD48 with a low affinity and binding does not require
 glycosylation of CD2. *Eur J Immunol* 23: 1373-7
- Weiss, A. and Littman, D. R. (1994) Signal transduction by lymphocyte antigen

receptors. *Cell* 76: 263-74

Vita *et al.*, 1998, "Novel miniproteins engineered by the transfer of active sites to small natural scaffolds." *Biopolymers* 47:93-100.

- 5 Weisshoff *et al.*, 1999, "Mimicry of beta II'-turns of proteins in cyclic pentapeptides with one and without D-amino acids." *Eur. J. Biochem.* 259:776-788.

Johannesson *et al.*, 1999, "Bicyclic tripeptide mimetics with reverse turn inducing properties." *J. Med. Chem.* 42:601-608.

Downloaded from www.jbc.org/